




Sören Starck (Autor)

Microbial production of *cis,cis*-muconic acid from hydrothermally converted lignocellulose



<https://cuvillier.de/de/shop/publications/8615>

Copyright:

Cuvillier Verlag, Inhaberin Annette Jentsch-Cuvillier, Nonnenstieg 8, 37075 Göttingen, Germany

Telefon: +49 (0)551 54724-0, E-Mail: info@cuvillier.de, Website: <https://cuvillier.de>

Table of Contents

| | |
|--|-------------|
| Summary | VIII |
| Zusammenfassung | IX |
| 1 Introduction | 1 |
| <u>1.1 General introduction</u> | 1 |
| <u>1.2 Objectives</u> | 2 |
| 2 Theoretical Background | 3 |
| <u>2.1 Biomass as a resource</u> | 3 |
| 2.1.1 Economic perspectives | 3 |
| 2.1.2 Lignin - a highly underexploited renewable | 4 |
| 2.1.3 Hemicellulose – a very abundant polymer..... | 11 |
| <u>2.2 Depolymerization of biomass</u> | 12 |
| 2.2.1 The pulping process | 12 |
| 2.2.2 Biochemical lignin depolymerization in nature..... | 15 |
| 2.2.3 Technical depolymerization of lignin..... | 16 |
| 2.2.4 Use of genetically engineered plants..... | 19 |
| <u>2.3 Bioconversion of biomass-derived carbon</u> | 20 |
| 2.3.1 <i>Pseudomonas putida</i> | 20 |
| 2.3.2 <i>Corynebacterium glutamicum</i> | 22 |
| 2.3.3 <i>Amycolatopsis sp.</i> | 25 |
| <u>2.4 Cis,cis-muconic acid - an important platform chemical</u> | 25 |

| | | |
|----------|---|-----------|
| 3 | Material and Methods | 29 |
| 3.1 | <u>Lignin and hemicellulose</u> | 29 |
| 3.2 | <u>Hydrothermal lignin conversion</u> | 30 |
| 3.3 | <u>Hydrothermal lignin conversion at miniaturized scale</u> | 31 |
| 3.4 | <u>Hydrothermal hemicellulose conversion</u> | 32 |
| 3.5 | <u>Fractionation of depolymerized hemicellulose by distillation</u> | 32 |
| 3.6 | <u>Distillation of lignin hydrolysate</u> | 33 |
| 3.7 | <u>Concentration of lignin and hemicellulose hydrolysates</u> | 33 |
| 3.8 | <u>Decolorization of hemicellulose hydrolysates</u> | 34 |
| 3.9 | <u>Microorganisms and plasmids</u> | 34 |
| 3.10 | <u>Media</u> | 34 |
| 3.11 | <u>Cultivation of <i>Pseudomonas putida</i></u> | 37 |
| 3.12 | <u>Cultivation of <i>Corynebacterium glutamicum</i> strains</u> | 38 |
| 3.13 | <u>Cultivation of <i>Amycolatopsis sp.</i></u> | 39 |
| 3.14 | <u>Fed-batch production of <i>cis,cis</i>-muconic acid</u> | 39 |
| 3.15 | <u>Quantification of glucose and xylose</u> | 40 |
| 3.16 | <u>Quantification of mono- and disaccharides</u> | 40 |
| 3.17 | <u>Quantification of aromatics and muconic acid derivatives</u> | 41 |
| 3.18 | <u>Fingerprinting of aromatics in lignin hydrolysates</u> | 41 |
| 3.19 | <u>Quantification of organic acids and alcohols</u> | 42 |
| 3.20 | <u>Quantification of cell concentration</u> | 42 |

| | | |
|----------|--|------------|
| 4 | Results and Discussion | 43 |
| 4.1 | <u>Microbial production of <i>cis,cis</i>-muconic acid from aromatics</u> | 43 |
| 4.2 | <u>Cascaded production of <i>cis,cis</i>-muconic acid from Kraft lignin</u> | 47 |
| 4.2.1 | Hydrothermal conversion of Kraft lignin | 47 |
| 4.2.2 | Production of <i>cis,cis</i> -muconic acid from lignin hydrolysate | 51 |
| 4.3 | <u>Impact of the lignin-type on depolymerization</u> | 54 |
| 4.3.1 | Screening of hydrothermal conversion efficiency | 54 |
| 4.3.2 | Optimized conversion of Kraft and organosolv lignin | 58 |
| 4.3.3 | Improvement of the aromatic | 59 |
| 4.3.4 | Bioconversion of lignin hydrolysates | 62 |
| 4.4 | <u>Gram-scale production of <i>cis,cis</i>-muconic</u> | 68 |
| 4.5 | <u>Use of hemicellulose-derived sugars as co-substrate</u> | 74 |
| 4.5.1 | Hemicellulose depolymerization into sugar monomers | 74 |
| 4.5.2 | Purification and vacuum concentration | 82 |
| 4.5.3 | <i>Cis,cis</i> -muconic acid production using xylose | 84 |
| 5 | Conclusion and Outlook | 92 |
| 6 | Appendix | 96 |
| 6.1 | <u>Abbreviations</u> | 96 |
| 6.2 | <u>Cultivation of <i>P. putida</i> MA-9 on organosolv lignin hydrolysate</u> | 97 |
| 6.3 | <u>Monomers obtained from multiple runs of HTC</u> | 98 |
| 6.4 | <u>Cultivation on hemicellulose hydrolysate</u> | 99 |
| 7 | References | 102 |